

Some recent progress in the weak noise theory of the KPZ equation

Yier Lin (University of Chicago)

November 14, 2022

Stochastic PDEs and Related Topics


Joint work with Pierre Yves Gaudreau Lamarre and Li-Cheng Tsai

The KPZ equation

- The KPZ equation is a stochastic PDE that models the random surface growth:

$$\partial_t h = \frac{1}{2} \partial_{xx} h + \frac{1}{2} (\partial_x h)^2 + \xi.$$

The growth subjects to **smoothing mechanism**, **slope dependence**, and **space-time independent noise**.



Freidlin-Wentzell Large Deviation Principle (LDP)

The theory of Freidlin-Wentzell studies the **large deviation principle (LDP)** of **small noise perturbation** of dynamical systems.

In our context, we consider the KPZ equation with a **small parameter** ε ,

$$\partial_t h_\varepsilon = \frac{1}{2} \partial_{xx} h_\varepsilon + \frac{1}{2} (\partial_x h_\varepsilon)^2 + \sqrt{\varepsilon} \xi.$$

The formalism is standard in stochastic analysis literature.

Short time formalism

The process $h_\varepsilon(t, x)$ has the same law as $h_1(\varepsilon^2 t, \varepsilon x)$.

The Freidlin-Wentzell LDP

The Freidlin-Wentzell LDP studies the **atypical behavior** of h_ε .


The typical behavior

As $\varepsilon \rightarrow 0$, $h_\varepsilon \rightarrow h_0$, where h_0 solves the HJ equation

$$\partial_t h_0 = \frac{1}{2} \partial_{xx} h_0 + \frac{1}{2} (\partial_x h_0)^2.$$

The atypical behavior (one-point)

Fix $(t, x) = (2, 0)$. Study the probability $\mathbb{P}(h_\varepsilon(2, 0) \approx \lambda)$, where $\lambda \neq h_0(2, 0)$.




pictures courtesy of Li-Cheng Tsai

The one-point LDP

The one-point LDP refers to, as $\varepsilon \rightarrow 0$,

$$\mathbb{P}(h_\varepsilon(2,0) \approx \lambda) \approx \exp\left(-\varepsilon^{-1} I_{op}(\lambda)\right),$$

where I_{op} is referred as the (one-point) **rate function**.



Precise meaning

More precisely, the above notation means

$$\lim_{\delta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \varepsilon \log \mathbb{P}(|h_\varepsilon(2,0) - \lambda| \leq \delta) = -I_{op}(\lambda).$$

The Freidlin-Wentzell LDP

The Freidlin-Wentzell (process level) LDP: For $g \in C([0, 2] \times \mathbb{R})$,

$$\mathbb{P}(h_\varepsilon \approx g) \approx e^{-\varepsilon^{-1}I(g)}.$$

Precise meaning

For any measurable set $\Omega \subseteq C([0, 2], \mathbb{R})$, we have

$$-\inf_{g \in \Omega^\circ} I(g) \leq \liminf_{\varepsilon \rightarrow 0} \varepsilon \log \mathbb{P}(h_\varepsilon \in \Omega) \leq \limsup_{\varepsilon \rightarrow 0} \varepsilon \log \mathbb{P}(h_\varepsilon \in \Omega) \leq -\inf_{g \in \overline{\Omega}} I(g).$$

Condition on $h_\varepsilon(2, 0) = \lambda$, we want study the $\varepsilon \rightarrow 0$ limit of h_ε , which is called **the most probable shape**.

The most probable shape is given by

$$\arg \min \left\{ I(g) : g(2, 0) = \lambda \right\}.$$

Different approaches

Exact formula approach

Use the exact formula to study the one-point LDP rate function.

Obtain the exact one-point rate function.

Krajenbrink, Le Doussal,
Majumdar, Rosso, Schehr, ...

Weak noise theory approach

Study the process-level LDP and most probable shapes.

Hartmann, Janas, Kamenev,
Katzav, Kolokolov, Kurshnov,
Meerson, Sasorov, ...

Gaudreau Lamarre, L, Tsai, ...

The weak noise theory + integrable system (2021+)

Bettelheim, Krajenbrink, Le Doussal, Meerson, Smith, ...

Tsai, ...

Theorem (L-Tsai 21)

Take $g \in C([0, 2] \times \mathbb{R})$, we have

$$\mathbb{P}(h_\varepsilon \approx g) \approx e^{-\varepsilon^{-1}I(g)},$$

the rate function I will be specified later.

The definition of I

It is known that for $\rho \in L^2([0, 2] \times \mathbb{R})$,

$$\mathbb{P}(\sqrt{\varepsilon}\xi \approx \rho) \approx e^{-\varepsilon^{-1} \cdot \frac{1}{2} \|\rho\|_{L^2}^2}.$$

We define $H[\rho]$, the solution to

$$\partial_t H = \frac{1}{2} \partial_{xx} H + \frac{1}{2} (\partial_x H)^2 + \rho.$$

The rate function is given by

$$I(g) = \inf \left\{ \frac{1}{2} \|\rho\|_{L^2}^2 : H[\rho] = g \right\}.$$

The minimizer of variational formula

By the expression of I , we have

$$\begin{aligned} & \inf \{ I(g) : g(2, 0) = \lambda \} \\ & = \inf \left\{ \frac{1}{2} \|\rho\|_{L^2}^2 : \mathbf{H}(\rho)(2, 0) = \lambda \right\} \end{aligned}$$

Proposition (Gaudreau Lamarre-L-Tsai 21)

The minimizer of the inf exists.

Question (Uniqueness?)

The minimizer of variational formula

By the expression of I , we have

$$\begin{aligned} & \inf \{ I(g) : g(2, 0) = \lambda \} \\ &= \inf \left\{ \frac{1}{2} \|\rho\|_{L^2}^2 : \mathbf{H}(\rho)(2, 0) = \lambda \right\}. \end{aligned}$$

Proposition (Gaudreau Lamarre-L-Tsai 21)

The minimizer of the inf exists.

Question (Uniqueness?)

*It depends on the initial condition and value of λ . For Brownian initial condition, a symmetry breaking (when λ is large) has been predicted by *Hartmann, Janas, Kamenev, Krajenbrink, Le Doussal, Meerson, Sasorov ...**

Narrow wedge initial data

The Hopf-Cole solution

The solution to the KPZ equation

$$\partial_t h = \frac{1}{2} \partial_{xx} h + \frac{1}{2} (\partial_x h)^2 + \xi$$

can be defined via $h = \log Z$, where Z solves the stochastic heat equation

$$\partial_t Z = \frac{1}{2} \partial_{xx} Z + \xi Z.$$

Narrow wedge initial data

We set $Z(0, \cdot) = \delta(\cdot)$. As $t \rightarrow 0$,

$$h(t, x) \approx -\frac{x^2}{2t} - \log \sqrt{2\pi t}.$$

Proposition (Tsai 22)

Under narrow wedge initial condition, the minimizer of

$$\inf \left\{ \frac{1}{2} \|\rho\|_{L^2}^2 : \mathbf{H}(\rho)(2, 0) = \lambda \right\}$$

is unique when $\lambda \in (-\infty, \lambda_0]$ for some λ_0 .

Define the unique minimizer as ρ_λ^m .

Let (w_λ, h_λ) be a scaled version of $(\rho_\lambda^m, \mathbf{H}(\rho_\lambda^m))$.


A lower-tail ($\lambda \rightarrow -\infty$) limit

Theorem (L-Tsai 22)

Uniformly over the compact domain in $[0, 2] \times \mathbb{R}$, we have

$$\lim_{\lambda \rightarrow -\infty} h_\lambda = h_*^-.$$

This confirms the prediction from Kamenev, Kolokolov, Korshunov, Meerson, Sasorov, ...



A lower-tail ($\lambda \rightarrow -\infty$) limit

The (w_λ, h_λ) satisfies a pair of PDEs. Taking $\lambda \rightarrow -\infty$ limit and solving the limiting equation, one gets

$$w_*(t, x) = -\frac{1}{2\pi\ell(t)} \left(1 - \frac{x^2}{\ell(t)^2}\right)_+,$$

where ℓ is implicitly defined by

$$|t - 1| = \sqrt{1 - \frac{\pi\ell(t)}{2}} + \frac{2}{\pi} \tan^{-1} \sqrt{\frac{2}{\pi\ell(t)} - 1}.$$

Proposition (L-Tsai 21)

As $\lambda \rightarrow -\infty$, $w_\lambda \rightarrow w_*$ in L^2 .

The lower-tail ($\lambda \rightarrow -\infty$) limit

By Feynman-Kac formula,


$$h_\lambda(t, x) = |\lambda|^{-1} \log \mathbb{E} \left[\exp \left(|\lambda| \int_0^t w_\lambda(s, W_{\lambda, x}(s)) ds \right) p_\lambda(t, x) \right],$$

where $W_{\lambda, x}(s)$ is a Brownian bridge with diffusive parameter $|\lambda|^{-1/2}$ and $p_\lambda(t, x) = \frac{1}{\sqrt{2\pi|\lambda|^{-1}}} e^{-\frac{|\lambda|x^2}{2t}}$.

Replacing w_λ with w_* and taking $\lambda \rightarrow -\infty$, we have

$$h_*^-(t, x) = \sup_\gamma \left(\int_0^t \underbrace{w_*(s, \gamma(s))}_{\text{energy}} - \underbrace{\frac{1}{2}\dot{\gamma}(s)^2}_{\text{entropy}} ds \right).$$

Maximize the combination of energy and entropy.




A class of geodesics

We call the maximizers as **geodesics**.

Proposition (L-Tsai 21)

For any $(t, x) \in (0, 2] \times \mathbb{R}$, the geodesics from (t, x) to $(0, 0)$ are as depicted in the figure.




Some remarks on the proof

Knowing the L^2 convergence $w_\lambda \rightarrow w_*$ is not enough. It is **not true** that for arbitrary $\tilde{w}_\lambda \rightarrow w_*$ in L^2 , we have $h_\lambda(\tilde{w}_\lambda) \rightarrow h_*$.
Extra ideas are needed.

The $\lambda \rightarrow \infty$ limit.

In [Gaudreau Lamarre-L-Tsai 21], we also obtain h_*^+ : The $\lambda \rightarrow \infty$ limit of most probable shape.

- Show that w_λ is nearly time-independent.
- Study the ground state of Schrödinger-type operator.
- Identify the ground state using Gagliardo-Nirenberg-Sobolev inequality.



Some more remarks


The $\lambda \rightarrow \pm\infty$ limit

- For $\lambda \rightarrow -\infty$, w_λ spans a wide region in spacetime.
- For $\lambda \rightarrow \infty$, w_λ vanishes everywhere except in a small neighborhood around $x = 0$.

Minimize $\|w_\lambda\|_{L^2}$ so that

$$\mathbb{E} \left[\exp \left(|\lambda| \int_0^t w_\lambda(t-s, W_\lambda(s)) ds \right) \right] = e^\lambda,$$

where $W_\lambda(s) = |\lambda|^{-1/2} B_b(s)$.



Generalize the theory for other initial data, for example Brownian?

Uniqueness of the minimizer and symmetry breaking.

...



Thank you!